

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## INDEX TO VOLUME VIII

New names and the final members of new combinations are in bold face type

```
Ambrosiaceae, 17
 Abies balsamea, 107; grandis, 91;
   lasiocarpa, 91; nobilis, 91
                                         Amelanchier Bakeri, 152; mormon-
 Acacia Greggii, 164
                                          ica, 152; oreophila, 146, 153; poly-
 Acer nigrum, 101
                                           carpa, 146
 Achlea Klebsiana, 108
                                         American mycological literature, In-
 Achlys triphylla, 102
                                          dex to, 59, 116, 188, 227, 289
 Adenocaulon bicolor, 103
                                         American species of Ascodesmis.
 Adiantum latifolium, 25
                                          North, 1
                                        Amorpha californica, 169; canescens,
 Aecial stage of Coliosporium ribi-
   cola, The, 309
                                           169; nana, 168
                                        Ampelomyces quisqualis, 175
 Aecidiaceae, 23
Aecidium, 131, 134; Abroniae, 150;
                                        Ampelopsis quinquefolia, 149
   Allenii, 150; Brandegei, 159; Cock-
                                        Amsonia, 135; salicifolia, 134
   erellii, 150; Compositarum, 150;
                                        Amvgdalaceae, 16
   Euphorbiae, 151; Eurotiae, 151;
                                        Amygdalus persica, 16
   Fendleri, 159; Hydrophylli, 138,
                                        Anastrophia bahamensis, 19
   141, 160; Iridis, 129; Lepidii, 163;
                                        Anchusa, 130; officinalis, 133, 141
  lini, 55; Nymphoidis, 16; obesum,
                                        Andira, 45
   134, 141; Phaceliae, 160; Physa-
                                        Andropogon brevifolius, 21; Schoe-
  lidis, 151; Sphaeralceae, 160, 161
                                          nanthus, 21
Agaricaceae, 251, 297
                                        Anemone, 128; canadensis, 128; cyl-
                                          indrica, 127, 128, 132
Agaricales, 172
                                        Angioridium sinuosum, 206; valva-
Agaricus arvensis, 70, 72; campes-
  ter, 172, 186, 297; discretus, 317;
                                          tum, 206
  echinocephalus, 232; onustus, 232;
                                        Anogra coronopifolia, 144
                                        Anthephora elegans, 19; hermaphro-
  placomyces, 297; ruber, 215
Agave neomexicana, 145
                                          dita, 19
Agrimonia striata, 164
                                        Anthurium, 22; scandens, 22
Agropyron, 138, 139; occidentale,
                                        Apios, 98; tuberosa, 98
  170; Smithii, 146, 161, 170; tene-
                                        Apocynum, 134; cannabinum,
                                          141; hypericifolium, 134
  rum, 138, 141
                                        Apple trees, Fungi producing heart-
Albugo Bliti, 144; candida, 144;
                                          rot of, 5
  Froelichiae, 144;
                      Ipomaeae-pan-
                                        Aquilegia, 128; caerulea, 157; cana-
  duratae, 144; Trianthemae, 145
                                          densis, 132
Aleuria repanda, 197
                                        Arabis Fendleri, 160; oxyphylla, 160
Allionia melanotricha, 150
                                        Araceae, 22
Althaea rosea, 159
                                        Arcyria bicolor, 210; cinerea, 41, 208,
Alveolaria Cordiae, 18
                                          210, 211; denudata, 41; globosa,
Amanita, 52, 53, 114, 231; Caesarea,
                                          210; incarnata, 41, 210; insignis,
  225, 251; Frostiana, 251; mus-
                                          41; minor, 210; pallida, 211
  caria, 114; nivalis, 233; onusta,
                                        Argomyces Oxalidis, 19, 151; Ver-
  299; pantherina, 114; phalloides,
                                          noniae, 24
  114, 233, 251; phalloides striatula.
                                        Aristida bromoides, 156
  232; rubescens, 251; solitaria, 231
                                        Arkville, New York, Fungi collected
Amanitopsis, 53; agglutinata, 251;
                                         at, 293
  albocreata, 232, 251; vaginata, 186
                                        Armillaria, 53, 183; evanescens, 53;
Amaranthus blitoides, 144; Powellii,
                                         mellea, 65, 114, 183
                                       Artemisia cana, 156; dracunculoides,
Amarella heterosepala, 166; stricti-
                                         146; franserioides, 163; nova, 156;
 flora, 166
                                         redolens, 156; silvicola, 156; tri-
Amauroderma Chaperi, 314
                                         dentata, 156
```

Basidiomycetes collected in Indo-Arthur, J. C., Cultures of Uredineae in 1915, 125; Uredinales of Porto China by C. B. Robinson, 214 Battarea laciniata, 174 Rico based on collections by F. L. Stevens,1 16 Belonidium Macounii, 98, 103 Artocarpaceae, 17 Berberis Fendleri, 159 Berry, Edward W., Remarkable fossil Artocarpus Camansi, 26; communis, fungi, 73 Betula populifolia, 105 Arundinaria, 128 Biatora phaeophora, 245 Asclepiadaceae, 16 Bibliography and new species of Phil-Asclepias, 134, 135; brachystephana, ippine fungi, 253 159; curassavica, 16; galioides, Bidens pilosa, 25; tenuisecta, 170 159; latifolia, 159; pulchra, 134; Bignoniaceae, 18 syriaca, 134, 135, 141; subverticil-Bihai psittacorum, 19 Bjerkandera adusta, 296 Ascobolus, 93, 94; albinus, 94, 95; Boletaceae, 249, 251, 296 atrofuscus, 95; caninus, 3; carbo-Boletinellus porosus, 296 narius, 93, 94, 95, 97; carbonicola, 95; geophilus, 93, 94, 96, 97; mi-Boletus, 225; auriporus, 251; felleus, 251; katui, 216; luteus, 53; reticucroscopicus, 3; subglobosus, 94, latus, 217; retipes, 251; rubritubi-96, 97; viridis, 93, 95, 97 Ascobolus, The earth-inhabiting spefer, 54; sanguineus, 215; Satanas, cies of, 93 114; tenuis, 217 Boudiera canina, 3; Claussenii, 3; Ascochyta, 104; Achlydis, 101; Pisi, f. Lupini, 104 microscopica, 3 Bouteloua, 134; aristidoides, 171; Ascodesmis, 1, 2; microscopica, 2, barbata, 171; breviseta, 171; cur-3, 4; nigricans, 1, 2, 3; porcina, tipendula, 159, 171; eriopoda, 171; 2, 3, 4; reticulata, 3 gracilis, 171; oligostachya, 171 Ascodesmis, North American species Bovista pila, 299 of, 1 Bradburya pubescens, 46 Ascomycetes, 295 Ascomycetes of Ohio, Fink's, 57 Brassica, 43; napi, 43; rapae, 43 Brenckle, J. F., Lamprospora detonia Asparagus officinalis, 156 sp. nov., 318 Asprella Hystrix, 138 Bromus polyanthus, 171 Aster, 130; laevis, 158; paniculatus, 130; vallicola, 146 Bruner, Stephen C., A new species of Asterina mexicana, 145 Endothia, 239 Bulbilis dactyloides, 171 Astragalus, 165; amphioxys, 165; Bulgaria bicolor, 237 Bigelovii, 165; bisulcatus, 170; Bullaria, 136; tumidipes, 136 165; crassicarpus, Drummondii, 170; missouriensis, 170; mollissi-Caesalpiniaceae, 16, 18, 20 mus, 165; Nuttallianus, 170; Thur-Cajanus indicus, 42 beri, 165; Wootoni, 165, 170 Calamovilfa longifolia, 126, 127 Attractive species of Melanoleuca Calapogonium orthocarpum, 45 from Oregon, An, 113 Calathea lutea, 25 Auerswaldia Pringlei, 146 Calliospora Farlowii, 18; Petaloste-Auricularia Auricula, 52, 56, 314; nimonis, 151 grescens, 314 Camarophyllus basidiosus, 297; Can-Avena fatua, 161; sativa, 171, 172 tharellus, 297; pallidus, 297 Camarosporium, 104; coronillae var. Baccharis, 18; glutinosa, 152 Spiraeae, 104; Spiraeae, 104 Baccilus Mangiferae, 223 Cantharellus cibarius, 251;

<sup>1</sup> This article contains an index to the Uredinales of the West Indies and also an index to their hosts.

Bacterium solanacearum, 109
Badhamia capsulifera, 206; decipiens,

206; rubiginosa, 201

205, 206; panicea, 35; papaveracea,

Based on collections by F. L. Stevens,

Uredinales of Porto Rico, 16

barinus, 251; infundibuliformis,

Carex, 169; arctata, 130, 141; Doug-

lasii, 156; durifolia, 131; Jamesii,

251; minor, 251

Carduaceae, 18, 19

Capriola Dactylon, 20

```
131; pubescens, 130; tenuis, 130,
                                          Chytridales, 144
                                          Ciboria nebulosa, 295
  141
Carya, 315
                                          Cineraria palustris, 131
                                          Cintractia Caricis, 169; leucoderma,
Cassava, 44
Cassia, 16; alata, 44; angustisiliqua,
                                            225, 226
  18; bauhinioides, 164; emarginata,
  16; occidentalis, 44
                                          Cirsium pallidum, 157; undulatum,
Castanea mollissima, 55
                                            157; Wrightii, 157
Castilleja confusa, 151; integra, 151;
                                          Citrus, 112
  linariaefolia, 151; sulphurescens,
                                          Cladium effusum, 19
                                          Cladoderris dendritica, 56, 314
  151
Ceanothus americanus, 100
                                          Cladosporites, 77, 78; fasciculatus,
Celtis reticulata, 149
                                            77, 79; oligocaenicum, 77, 79
Cenangium populneum, 302, 306
                                          Cladosporium, 77; herbarum, 78, 176
Cenchrus carolinianus, 25
                                          Cladothrix lanuginosa, 144
Cephalanthus, 134; occidentalis, 134
                                          Clastoderma Debaryanum, 39
Ceratiomyxa fruticulosa, 35; fruticu-
                                          Claudopus nidulans, 173; subnidu-
  losa flexuosa, 35
                                            lans, 313
Cercospora, 42;
                   Amaryllidis,
                                          Clavaria cristata, 295; fusiformis,
  Amoraciae, 43; argythamniae, 55;
Bloxami, 43; borinquensis, 45;
                                            250, 294, 295; Kunzei, 295; pistil-
                                            laris, 53
  Bradburyae, 46; brassicola, 43;
                                          Clavariaceae, 250
  Cajani, 42; Cassavae, 44; Chamae-
                                          Claviceps purpurea, 146
  cristae, 44; conspicua, 42; densis-
                                          Clematis ligusticifolia, 128, 132, 157,
  sima, 44; guanicensis, 45; Hi-
bisci, 44; Hydropiperis, 43; Lepi-
dii, 43; Malachrae, 45; mari-
                                         Cleome, 42
                                          Clitocybe, 68, 317; adirondackensis,
  caoensis, 44; Mucunae, 43; namae, 55; Phyllitidis, 44; Polygo-
                                           297; candida, 69; eccentrica, 297; flaccida, 185; illudens, 251; infun-
  norum, 43; portoricensis, 42; rosi-
                                            dibuliformis, 251, 297; laccata,
                                            251; lactariiformis, 297; mona-
  cola, 43; Stevensii, 45; Vaginae,
                                            delpha, 68; nebularis, 65-69, 71;
Ceriomyces bicolor, 296; communis,
                                            nobilis, 69; robusta, 69
  296; illudens, 296; retipes, 296;
                                          Clitopilus abortivus, 65
  viscidus, 53, 296
                                          Coffea arabica, 16
Ceropteris triangularis, 153
                                         Coleosporium Eupatorii, 18; ribicola,
Cerrena unicolor, 296
                                            151, 309
Chaetochloa imberbis, 25
                                         Coleosporium ribicola, The aecial
                                            stage of, 309
Chamaecrista Aeschynomene, 20
Chamaecyparis lawsoniana, 92
                                         Collections by F. L. Stevens, Uredi-
                                           nales of Porto Rico based on, 16
Chamaenerion angustifolium, 164
Chamaesaracha Coronopus, 150
                                         Collybia agricola, 218; avellanei-
                                            disca, 218; avellaneigrisea, 218;
Chamaesyce albomarginata, 168; Fend-
  leri, 168; Geyeri, 168; glyptosper-
                                           badiialba, 218; cinchonensis, 218;
  ma, 168; hypericifolia, 25; lata,
                                            cremeimellea, 218; densifolia,
  168; petaloidea, 168; serpens, 168;
                                           218; dentata, 218; denticulata,
  serpyllifolia, 167; stictospora, 168
                                           218; domestica, 218; Earleae,
                                           218; Eatonae, 218; farinacea, 218;
Chanterel Chantarellus, 297; flocco-
                                           fimetaria, 219; flavescens, 219;
  sus, 297; infundibuliformis, 297
Cheilanthes Pringlei, 153
                                           fulvidisea, 219; fulvipes, 219;
Chenopodium, 129; album, 129, 135,
                                           Glatfelteri, 219; griseifolia, 219;
  141, 162, 176; Fremonti, 163; pra-
                                           jamaicensis, 219; ludoviciana,
                                           219; marasmiiformis, 219; mon-
  tericola, 163
Chlamydopus clavatus, 174
                                           ticola, 219; musicola, 219; ni-
                                           gritiformis, 219; oculata, 219;
Chloris elegans, 129, 165
Chlorosplenium aeruginosum, 56
                                           orizabensis, 219; pallida, 219;
Chondrioderma
                   subdictyospermum,
                                           platyphylla, 251; radicata, 251; ro-
                                           seilivida, 219; setulosa, 219; sinuata. 219; squamiger, 219;
Chrysanthemum leucanthemum, 104
Chrysothamnus linifolius, 146; Va-
                                           subavellanea, 219; subflaves-
  seyi, 146
                                           eens, 219; subflavifolia, 219;
```

326 Mycologia

sublatericia, 219; subnivulosa, Cudonia lutea, 295 Cultures of Uredineae in 1915, 125 219; subrugosa, 219; tenuifolia, 219; tortipes, 219; trullisata, 219; unakensis, 219; virgin-Cup-fungi—III. Peziza domiciliana and Peziza repanda, 195; IV. Peiana, 219; Volkertii, 219; xuchiziza clypeata, Photographs and delensis, 219 scriptions of, 235 Notes on the Coltricia cinnamomea, 296; peren-Curtis Herbarium, nis, 296 Myxomycetes of the, 199 Colus Schellenbergiae, 183; java-Cyathia hirsuta, 174 nicus, 184 Cycloporellus iodinus, 314 Cylindrosporium, 106; Chrysanthemi, Colus from Pennsylvania, A new 104; Crataegi var. brevispina, species of, 183 105; spigeliae, 55; Toxicodendri, Comandra pallida, 152 Comatricha nigra, 39; pulchella, 209; 105 Cynodon Dactylon, 21 typhoides, 39; typhoides var. heterospora, 39 Cyperaceae, 16, 17, 19 Conifers, Laboratory tests on the Cyperus, 225; distans, 25; esculentus, durability of American woods-I. 157; Gatesii, 225; odoratus, 25 Cyphella porrigens, 53 Flask tests on, 80 Convolvulcaeae, 19 Cytodiplospora parallela, 101 Coprinus, 53, 54, 70, 71, 186; atra-Cytospora chrysosperma, 295 mentarius, 70, 72, 173; atramentarius silvestris, 297; bisporus, 54; Daedalea, 216; amanitoides, 56, 216; applanata, 216; confragosa, 296; comatus, 52, 53; ephemerus, 186; flavida, 216; Palisoti, 216; philip-Hansenii, 54; micaceus, 70, 173, 186, 297; stercorarius, 186 pinensis, 110; repanda, 216 Cordia, 17; alliodora, 17; cylindro-Daldinia concentrica, 295 stachva, 18 Dangerous mushroom, A very, 186 Darluca filum, 175 Coriolopsis caperata, 314; occiden-Dasylirion Wheeleri, 148 talis, 56 Coriolus abietinus, 173, 296; haedi-Dasyochloa pulchella, 171 nus, 56; limitatus, 173; membrana-Dasyscypha arida, 53 ceus, 56; nigromarginatus, 296; Dearness, John, New or noteworthy pinsitus, 56; pubescens, 53; sector, species of fungi, 98 Delphinium Sapellonis, 157 56; versicolor, 173, 296 Cornus Nuttallii, 101 Descriptions of cup-fungi-III. Pe-Corticium pezizoideum, 53; serum, ziza domiciliana and Peziza re-184; varians, 184 panda, 195; IV. Peziza clypeata, Photographs and, 235 Cortinarius, 316; armillatus, 297; Diachaea leucopoda, 38; leucopoda chrysolitus, 54; semisanguineus, 297 Cortinellus vaccinus, 53 var. globosa, 38 Crataegus brevispina, 105 Dianthus caryophyllus, 181 Diaporthe, 100; columbiensis, 100; Euonymi, 99; Macounii, 100 Craterellus Cantharellus, 250, 295; cornucopioides, 250, 295 Craterium floriforme, 211, 213; leu-Dicaeoma poculiforme, 181 Dichrophyllum marginatum, 168 cocephalum var. cylindricum, 207, Dictydiaethalium plumbeum, 40, 211, 213; minimum, 207, 213; porphyrium, 207 213 Dictydium, 40; cancellatum, 40 Crepidopus, 218; ostreatus, 297 Cressa truxillensis, 157 Diderma, 38; antarctica, 37; hemi-Cribaria, 40; argillacea, 41, 212; elesphericum, 36 niveum, 36; niveum gans, 210; macrocarpa, 40; micro-Lyallii, 36; pallidum, 206; simplex, 37; spumarioides, 36; testaceum, scopica, 210; minima, 210; minutissima, 210; pyriformis, 40; splen-36; Trevelyani, 37 dens, 40 Didymaria Clematidis, 175 Didymium, 202; chrysopeplum, 200; Crinipellis fragilis, 110 Cronartium, 314; coleosporioides, 151; Clavus, 38; columbinum, 200, 209, 213; curtisii, 201; dealbatum, 201; Comandrae, 152; ribicola, 314 Crucibulum vulgare, 299 difforme, 212; eximium, 201, 202; Cryptogamme Stelleri, 153 luteogriseum, 201; megalosporum, Cryptosporium falcatum, 107 201, 202, 209, 213; melanospermum,

38; nectriaeforme, 202; nigripes, 201; nigripes var. eximium, 205, 213; obrusseum, 202; pusillum, 202; proximum, 202; pruinosum, 202; radiatum, 203; Ravenelii, 203; squamulosum, 203; terrigenum, 204, 213; tenerrimum, 203; xanthopus, 202 Ceanothi. Didymosphaeria 100; Housei, 100 Dimerosporium Collinsii, 146 Diplodia constricta, 102; Nut-taliae, 102; Spiraea, 102; Ulmi, Dirca, 135, 136; palustris, 135 Discina adnata, 237; clypeata, 237; orbicularis, 237; repanda, 198 Discomycetes, 98 Disella hederacea, 159, 161, 162; lepidota, 159 Distichlis maritima, 136; spicata, 135, 141, 163, 172 Dodge, B. O., Fungi producing heartrot of apple trees, 5 Dothichiza populea, 300-308 Dothichiza populea in the United States, 300 Dothidella, 45 Dothiorella, 111 Draba aurea, 160 Dugaldea Hoopesii, 148, 150, 176 Dulichium arundinaceum, 130, 140 Duportella Raimundoi, 110; velutina, 110 Durability of American woods-I. Flask tests on conifers, Laboratory tests on the, 80

Earlea speciosa, 152 Earliella corrugata, 314 Earth-inhabiting species of Ascobolus, The, 93 Ehretiaceae, 17, 18 Eichleriella, 316; gelatinosa, 316; Schrenkii, 316 Eleagnus, 128; argentea, 128 Eleocharis capitata, 25; flaccida, 25; mutata, 25 Elfvingia fomentaria, 296; megaloma, 296; tornata, 56, 314 Elfvingiella fasciata, 56, 314 Elmerina foliacea, 110 Elymus, 139; canadensis, 146, 157, 170; condensatus, 138; virginicus, 138, 141 239, 240; havanensis, Endothia, 241; longirostris, 242; parasitica, 112, 304; radicalis, 113, 239, 242 Endothia, A new species of, 239 Enerthenema papillatum, 39 Enteridium cinereum, 204

Entoloma lividum, 114, 123 Entyloma australe, 169; Compositarum, 170 Ephedra Torreyana, 155; trifurca, 154, 155 Epidendrum difforme, 21; rigidum, Epilobium adenocladon, 158; novomexicanum, 164 Erigeron subtrinervis, 146 Eriogonum cernuum, 166; effusum, 166; lonchophyllum, 166; racemosum, 166 Eriophorum, 131; angustifolium, 131; polystachion, 132; tenellum, 132; virginicum, 132; viridi-carinatum, 131, 132, 141 Eriosporangium evadens, 18; punctato-striatum, 152 Erysiphe, 147; Cichoracearum, 146-148, 175; Galeopsidis, 147; graminis, 147; Polygoni, 147, 148 Eucalyptus, 239; botryoides, 240, 242; microphylla, 242; occidentalis, 242; robusta, 242; rostrata, 242 Euonymus atropurpureus, 99 Eupatorium macrophyllum, 18 Euphorbia corollata, 127 Euphorbiaceae, 24 Eurotia lanata, 151 Exoascales, 316 Exobasidium, 316 Exogonium arenarium, 19 Experiment with Panaeolus papilionaceus, An, 317

Fabaceae, 16, 18 Fagus antarctica, 38 Family of Hymenomycetes, A new, 56 Favolus variegatus, 56 Fendlera falcata, 152; rupicola, 152 Festuca, 133; Thuberi, 132, 141 Fimbristylis spadicea, 17 Bruce, Fink. Hermann Hasse,-Lichenist, 243 Fink's Ascomycets of Ohio, 57 Fistulina hepatica, 251 Flask tests on conifers, Laboratory tests on the durability of American woods-I, 80 Fomes, 214; annosus, 111; applanatus, 250; Auberianus, 314; Ellisianus, 173; fomentarius, 225; igniarius, 6; lobatus, 250; pinicola, 111; populinus, 296; rimosus, 214, 250; ungulatus, 296 Fomitella supina, 56, 314 Fossil fungi, Remarkable, 73 Frasera macrophylla, 168; speciosa, 168

328 Mycologia

Fuligo cinerea, 204, 213; septica, 36
Fulvifomes extensus, 314
Fungi—XXIII, 121; XXIV, 191;
XXV, 231; Illustrations of
Fungi Collected at Arkville, New
York, 293
Fungi, New or noteworthy species
of, 98
Fungi of New Mexico, 142
Fungi producing heart-rot of apple
trees, 5
Fungi, Remarkable fossil, 73
Fungi—II, Studies in Porto Rican
parasitic, 42

Galactinia succosa, 295 Galactopus, 218; succosus, 297 Galera Hypnorum, 297; tener, 297 Galium asperrimum, 161 Ganoderma Bakeri, 110; plicatum, 110; Tsugae, 296 Ganolobus, 202 Gasteromycetes, 299 Gaura coccinea, 167; glabra, 167; induta, 167; parviflora, 167; villosa, Gayophitum ramosissimum, 158 Geaster floriformis, 174; hygrometricus, 174; saccatus, 299 Genus of resupinate polypores, A new, 56 Geopetalum, 218, 297; albescens, 218; geophilum, 218; subelatinum, 218; subhaedinum, 218; tremelliforme, 218 Geopyxis amplispora, 198; pallidula, 197; vulcanalis, 53 Gloeophyllum hirsutum, 296 Gloeosporium, 106; lunatum, 223; Toxicodendri, 105, 106 Glycyrrhiza lepidota, 147, 166 Graff, Paul W., Basidiomycetes collected in Indo-China by C. B. Robinson, 214; Bibliography and new species of Philippine fungi,2 253 Grifola Berkeleyi, 296 Grindelia, 158; aphanactis, 148, 158, 177; subalpina, 158 Grossularia leptantha, 151, 158 Guepinia spathularia, 314 Guilandina crista, 45 Guinardia Aesculi, 224 Gutierrezia, 158; tenuis, 158 Gymnopus, 218; agricola. 218; avellaneidiscus, 218; avellaneigriseus, 218; badiialbus, 218; chrysopeplus, 297; cinchonensis, 218; cremeimelleus, 218; densifolius, 218; dentatus, 218; denticulatus, 218;

domesticus, 218; dryophilus, 297; Earleae, 218; Eatonae, 218; farinaceus, 218; fimetarius, 219; flavescens, 219; fulvidiscus, 219; fulvipes, 219; Glatfelteri, 219; griseifolius, 219; jamaicensis, 219; ludovicianus, 219; marasmiiformis, 219; monticola, 219; musicola, 219; myriadophyllus, 297; nigritiformis, 219; oculatus, 219; oculus, 297; orizabensis, 219; pallidus, 219; platyphyllus, 297; radicatus, 297; roseilividus, 219; setulosus, 219; sinuatus, 219; squamiger, 219; subavellaneus, 219; subflavescens, 219; subflavifolius, 219; sublatericius, 219; subnivulosus, 219; subrugosus, 219; tenuifolius, 219; tortipes, 219; trullisatus, 219; unakensis, 219; virginianus, 2 Volkertii, 219; xuchilensis, 219 Gymnosporangium, 313; bermudianum, 18; gracilens, 152; inconspicuum, 152; juvenescens, 152; koreaense, 222; Nelsoni, 153 Gyrocephalus rufus, 295 Gyroporus castaneus, 296; cyanescens, 296

Habenaria maculosa, 18 Halerpestes Cymbalaria, 157 Hamelia erecta, 23 Hapalopilus gilvus, 56; rutilans, 296 Harper, Edward T., Two parasitic mushrooms, 65 Hasse,-Lichenist, Hermann Edward, Heart-rot of apple trees, Fungi producing, 5 Hebeloma peckii, 55 Hedgcock, George G., and Hunt, N. Rex, Dothichiza populea in the United States, 300 Helianthus annuus, 158; ciliaris, 159; fascicularis, 149 Heliconia psittacorum, 19 Heliopsis scabra, 146 Helminthosporium Hydropiperis, 43 Helotium citrinum, 295 Helvella crispa, 53 Hemileia vastatrix, 16 Hemitrichia clavata, 41; Vesparium, 208, 211 Hermann Edward Hasse,-Lichenist, 243 Herpotrichia nigra, 53 Heteropogon contortus, 170 Hevea, 315

Hexagona alveolaris, 296; daedalea,

<sup>&</sup>lt;sup>2</sup> This article contains an index to new genera and species.

296 Hexagonia, 217; tenuis, 217 Hiatula, 317; Benzonii, 317; ciliatula, 317; discreta, 317; minima, 317; purpurascens, 317 Hibiscus tiliaceus, 44 Hilaria Jamesii, 172; mutica, 172 Holcus jalapensis, 16; lanatus, 128; Sorghum, 16 Hordeum jubatum, 128, 161, 167; nodosum, 140; pusillum, 139-141 Host, A known species of smut on a new, 225 Humphrey, C. J., Laboratory tests on the durability of American woods —I. Flask tests on conifers, 80 Humulus Lupulus neomexicanus, 148 Hunt, Rex N., and Hedgcock, George G., Dothichiza populea in the United States, 300 Hyalopsora Cheilanthis, 153 Hydnaceae, 250, 296 Hydnum adustum, 250; erinaceum, 250; imbricatum, 53; repandum, 296; zonatum, 250 Hydrocotyle australe, 129 Hydrocybe ceracea, 297; flammea, 297; laeta, 297; miniata, 297; nitida, 297; Peckii, 297; punicea, 297 Hydrophyllum, 138; capitatum, 138, 141; Fendleri, 146, 160 Hygrophorus, 294, 317; miniatus, 251 Hymenocallis, 44 Hymenochaete pavonia, 110 Hymenoclea monogyra, 162 Hymenomycetes, 295 Hymenomycetes, A new family of, 56 Hypholoma appendiculatum, 298; incertum, 251; pecosense, 173 Hypochnus subtilis, 186 Hypocrea Richardsoni, 53 Hypomyces apiculata, 295; hyalinus, 295, 299; lactifluorum, 295 Hypoxis decumbens, 22 Hystrix, 138 Illustrations of fungi-XXIII, 121, XXIV, 191, XXV, 231 Illustrations of fungi I-XXII, Index to, 47 Index to American mycological literature, 59, 116, 188, 227, 289, 319 Index to illustrations of fungi I-XXII, 47 Indo-China by C. B. Robinson, Basidiomyctes collected in, 214 Inocybe, 298, 312, 316; anomala, 312 Insects and mushrooms, 113 Ionoxalis Martiana, 19; violacea, 151

56; lachnochaeta, 110; striatula,

Ipomoea arenaria, 19; hirsutula, 144; Steudeli, 19 Iris, 129; versicolor, 129 Irpex lacteus, 5 Irpiciporus lacteus, 296 Isocoma heterophylla, 163 Iva axillaris, 159; xanthiifolia, 146 Johnston, J. R., Marasmius on sugar cane, 115 Juglans major, 177 Juneus interior, 168; longistylis, 168 Juniperaceae, 18 Juniperus, 174; bermudiana, 18; californica, 91; chinensis, 222; lucayana, 18; monosperma, 152, 153; pachyphloea, 151; occidentalis, 91; scopulorum, 153; utahensis, 152, Kentucky fungi, Some, 249 Kniep's Beiträge zur Kenntnis der Hymenomyceten, III, Review of, Known species of smut on a new host, A, 225 Koeleria cristata, 160 Laboratory tests on the durability of American woods-I. Flask tests on conifers, 80 Laccaria laccata, 298; ochropurpurea, 315; striatula, 298 Lachnea lusatiae, 295; scutellata, 295 Lachnobolus cinereus, 204, 213 Lachnocladium Micheneri, 295; Schweinitzii, 295 Lactaria, 191; atroviridis, 191; cilicioides, 194; cinerea, 298; deliciosa, 114, 225; distans, 192; Gerardii, 251; Hibbardae, 298; hygrophoroides, 192, 298; lactiflua, 192, 251, 298; lignyota, 298; luteola, 192; maculosa, 193; piperata, 122, 185, 298; rufa, 192; subdulcis, 251, 298; subvelutina, 192; testacea, 192; torminosa, 114, 193; trivialis, 252; zonaria, 114 Lactuca integrata, 146; pulchella, 159 Laestadia biennis, 98; galactina, 55 Lamiaceae, 24 Lamproderma columbinum, 208 Lamprospora detonia, 318; trachycarpa, 318 Lapsana communis, 107 Larix europaea, 91 Lasiosphaeria hispida, 295 Lathyrus arizonicus, 166; decaphyllus, 166, 176; leucanthus, 147, 166 Laurinoxylon, 79

Leangium, 38

Lecanora redimita, 245 rans, 299; subincarnatum, 299 Lentinus, 217, 315; carneotomento-Lycopus, 131; virginicus, 181 sus, 298; crinitus, 56; exilis, 217; infundibuliformis, 315; lepideus, 11, Macrophoma celtidicola, 55; peck-82, 83; similis, 315; strigosus, 56, iana, 55 314 Macropodia macropus, 295; pubida, Lentodiellum, 218 295 Lenzites applanata, 216; betulina, Macrosporium Solani, 177 296; lurida, 217; pallida, 217; pa-Madia glomerata, 146 lisota, 216; platypoda, 217; repanda, Malachra rotundifolia, 46 216; subconfragosa, 217 Malvaceae, 16 Leocarpus fragilis, 36; lubrica, 295 Malvastrum elatum, 162 Leotia chlorocephala, 250 Mangifera indica, 239, 242 128; argentea, Lepargyraea. 173; Marantaceae, 25 canadensis, 128, 150, 177 Marasmius, 54, 115, 295; alliatus, Lepidium, 43; alyssoides, 163 298; caryophylleus, 54, 298; confluens, 298; delectans, 298; di-chrous, 298; foetidus, 298; gla-Lepidodendron, 75 Lepidostrobus, 75 bellus, 298; plicatus, 115; resino-Lepiota, 53, 54, 234; acutesquamosa, 233; aspera, 233; asperula, 233; sus, 298; rotula, 298; Sacchari, Cortinarius, 54; cristata, 313; erio-115; semiustis, 115; siccus, 252, phora, 233; Friesii, 233; Morgani, 298; stenophyllus, 115; subnudus, 252; seminuda, 313 298 Leptomyces, 218, 317 Marasmius on sugar cane, 115 Leptoporus armatus, 110; Bakeri, 110 Margarita metallica, 41 Leptostromella conigena, 105 Mariscus jamaicense, 19 Leucocrinum, 127 Marsonia, 106 Leucoporus ameides, 110 Marssonina Toxicodendri, 105 Leucosyris spinosus, 165 Medicago sativa, 150, 167; denticu-Levine, Michael, Review of Kniep's lata, 18; lupulina, 167 Beiträge zur Kenntnis der Hyme-Melampsora albertensis, 153; arctica, nomyceten, III, 184 153; Bigelovii, 153; Lini, 154; Licea, 37; antarctica, 37; applanata, Medusae, 154 211; artocreas, 211; macrospora, Melampsorella elatina, 154 212; microsperma, 212, 213; rubi-Melampsoropsis Pyrolae, 154 formis, 212; spermoides, 212; stipi-Melanconium parvulum, 105 tata, 212 Melanoleuca albissima, 298, 299; al-Lichenist, Hermann Edward Hasse, boflavida, 298; anomala, 312; melaleuca, 53, 298; olivaceiflava, 113 243 Melanoleuca from Oregon, An at-Ligularia sibirica, 131 tractive species of, 113 Ligusticum Porteri, 154 Melanopsamma waghornei, 55 Limnanthemum Grayanum, 17 Melanopus, 10 Limonium limbatum, 167 Meliola, 314 Mentha Penardi, 160 Lindbladia effusa var. simplex, 212 Linnaea, 149; americana, 149 Linum Lewisii, 154 Menyanthaceae, 17 Mertensia, 175; pratensis, 147 Metasphaeria **Macounii**, 100 Literature, Index to American mycological, 59, 116, 188, 227, 289 Lithospermum, 139 Long, W. H., Note on western red Microglossum rufum, 295 Micromphale, 218; badium, 219; fulrot in Pinus ponderosa, 178; The vifibrillosum, 219; subexcavatum, aecial stage of Coliosporium ribi-Microsphaera Symphoricarpi, 148 cola, 309 Lupinus aduncus, 175; ammophilus, Microstroma album, 177; Juglandis, 167; Kingii, 167; perennis, 104 177 Lycium pallidum, 136, 163, 141; vul-Mikania scandens, 19 Mollisia, 98; apiophila, 98 gare, 136, 141 Monarda stricta, 160 Lycogala Epidendrum, 41; flavo-Moneses uniflora, 154 fuscum, 41 Lycoperdales, 174 Moniliales, 176 Lycoperdon gemmatum, 299; sepa-Morchella crassipes, 53

Mucronoporus Andersoni, 56, 57 Nemopanthes mucronata, 107 Mucuna pruriens, 43 Neotiella albocinta, 295 Muhlenbergia neomexicana, 148; Por-Nephlyctis transformans, 18 teri, 172; Pringlei, 172; repens, Neurolaena lobata, 25 160; squarrosa, 161 New family of Hymenomycetes, A, 56 Murrill, W. A., An attractive species New genus of resupinate polypores, of Melanoleuca from Oregon, 113; Fungi Collected at Arkville, New A, 56 New Mexico, Fungi of, 142 York, 293; Illustrations of fungi-New or noteworthy species of fungi, XXIII, 121; XXIV, 191; XXV, 231; Index to illustrations of fungi 98 New species of Colus from Pennsyl-I-XXII, 47; A new family of Hyvania, A, 183 menomycetes, 56; A new genus of New species of Endothia, A, 239 New species of Philippine fungi, Bibresupinate polypores, 56; Pleurotus, Omphalia, Mycena, and Colliography and, 253 lybia published in North American Nidularia pisiformis, 299 Flora, 218; A very dangerous Nigredo Caladii, 181; caryophyllina, mushroom, 186 181; Medicaginis, 18; Trifolii, 18 Nolina microcarpa, 170 North American Flora, Pleurotus, Mushroom, A very dangerous, 186 Mushrooms, Insects and, 113 Mushrooms, Two parasitic, 65 Omphalia, Mycena, and Collybia Mycena, 53; Abramsii, 220; adipublished in, 218 rondackensis, 220; alcalini-North American species of Ascoformis, 220; argillacea, 220; desmis, r atribrunnea, 220; atridisca, 220; Notes on western red rot in Pinus aurantiaca, 220; aurantiidisca, ponderosa, 178 220; avellanea, 220; avellaneigrisea, 220; brevipes, 220; Notes and brief articles, 52, 108, 183, 222, 312 caesiialba, 220; carbonicola, 220; cervinialba, 220; cincho-Notes on the Myxomycetes of the Curtis Herbarium, 199 nensis, 220; cinereiavellanea, Noteworthy species of fungi, New or, 220; collybiiformis, 220; farinacea, 220; flavicitrina, 220; fuliginosa, 220; fumosiavel-Nothoholcus lanatus, 128 Notholaena, 153; sinuata, 153 lanea, 220; fusipes, 220; gracil-Nothoscordium, 139, 140; bivalve, lipes, 220; Grantii, 220; lateri-139-141; striatum, 139, 140 cia, 220; lepiotiformis, 220; Nuttallia cerasiformis, 100, 102 leucophaea, 220; longipes, 220; Nyssopsora echinata, 154 ludoviciana, 220; magna, 220; margarita, 220; melleidisca, Octospora varia, 197 221; minutissima, 221; murina, Odostemon Fremontii, 169; haema-221; myceliosa, 221; niveipes, tocarpus, 169; repens, 169 221; occidentalis, 221; ochra-Oenothera biennis, 167; procera, 148 ceicinerea, 221; paludicola, 221; Ohio, Fink's Ascomycetes of, 57 Ohleriella neomexicana, 148 parvula, 221; pectinata, 221; plumbeibrunnea, 221; pubes-Oidium fusisporioides f. Lapsanae, cens, 221; roseipallens, 221; ro-107 seola, 221; rutilantiformis, 221; Olyra, 22 Sabali, 221; scabripes, Omphalia acuminata, 219; Bakeri, subfumosa, 221; subpulveru-lenta, 221; subtenuipes, 221; 220; californiensis, 220; coccinea, 219; collybiiformis, 219; syringea, 221; tenuicula, 221; convexa, 220; cremea, testacea, 221; trojana, 221; vircuspidatella, 219; cuticolor, 220; idigrisea, 221 Dawsonii, 219; distantifolia, 220; Earlei, 219; hypobrunnea, Mycological literature, Index to American, 59, 116, 188, 227, 289 219; incarnata, 219; jalapensis, Myxomycetes, 199, 200 219; lenta, 219; luteicolor, 219; Myxomycetes from South America, McMurphyi, 220; miniata, 219; myceniformis, 220; niveicolor, Myxomycetes of the Curtis Herbar-219; petasiformis, 220; praedeium, Notes on the, 199 currens, 220; pseudogrisea, 220;

roriduliformis, 220; Sequoi-Pappophorum Wrightii, 171 arum, 220; subavellanea, 220; Parasitic fungi-II, Studies in Porto subcartilaginea, 220; subim-Rican, 42 maculata, 220; subscyphoides, Parasitic mushrooms, Two, 65 220; tepeitensis, 220; translu-Parmelia subolivacea, 245 centipes, 220; turbinata, 220 Parosela domingensis, 18 Omphalina, 218; acuminata, 219; Parthenocissus vitacea, 149 chrysophylla, 298; coccinea, 219; Paspalum fimbriatum, 25 Pavonia racemosa, 16 collybiiformis, 219; cremea, 219; cuspidatella, 219; Dawsonii, 219; Paxillus corrugatus, 252; involutus, Earlei, 219; hypobrunnea, 219; 298; Lepista, 69 incarnata, 219; jalapensis, 219; Pellaea andromedaefolia, 153 lenta, 219; luteicolor, 219; miniata, Peltandra virginica, 181 219; niveicolor, 219; Sequoiarum, Peniophora cinerea, 295 220; subcartilaginea, 220; sub-Pennsylvania, A new species of Colus scyphoides, 220; tepeitensis, 220 Omphalopsis, 218; Bakeri, 220; califrom, 183 Peperomia, 23; hernandiflora, 23 forniensis, 220; campanella, 298; Perichaena, 211; chrysosperma, 207; convexa, 220; cuticolor, 220; disdepressa, 211; vermicularis, 206, tantifolia, 220; fibula, 298; Mc-Murphyi, 220; myceniformis, 220; Peridermium, 314; coloradense, 154; petasiformis, 220; praedecurrens, Ephedrae, 154; ribicola, 309-311 220; pseudogrisea, 220; roriduli-Peronospora, 74; Lepidii, 145; paraformis, 220; subavellanea, 220; sitica, 145 Peronosporales, 144 subimmaculata, 220; translucentipes, 220; turbinata, 220 Peronosporites antiquarius, 75, 76 Peronosporoides, 76; palmi, 74, Onosmodium, 139 Ophiotheca pallida, 207; umbrina, 75, 77, 79 207; wrightii, 207 Persea gratissima, 239, 242 Oplismenus hirtellus, 21 Petalostemum oligophyllum, 151, 169; Orchidaceae, 18, 21 purpureum, 169 Oregon, An attractive species of Peziza, 236; Adae, 197; adnata, 235, Melanoleuca from, 113 237; amplispora, 197; Barlaeana, Oreobatus, 155; deliciosus, 155; neo-237; clypeata, 235, 236, 237; domimexicanus, 155 ciliana, 195, 196, 197; pallidula, Ornithogalum bivalve, 139; umbella-197; odorata, 197; orbicularis, 236, 237; regalis, 150; repanda, 195, tum, 139 Oryzopsis hymenioides, 172 196, 197, 295; repanda amplispora, Ostracoderma spadiceum, 213 197; Stevensoniana, 197; varia f. Otidea alutacea, 295; Auricula, 53 lignicola, 198; varia f. typica, 197 Overholts, L. O., and Mae F., Some Pezizales, 150 Kentucky fungi, 249 Phacelia, 139; heterophylla, 147, 160. Oxalidaceae, 19 175 Oxytropis Lamberti, 165 Phaeopezia Nuttallii, 95 Phaseolus vulgaris, 165 Pachyella Barlaeana, 237 Philadelphus argyrocalyx, 152; ellip-Pachylophus hirsutus, 167; macroticus, 152; microphyllus, 152 Philibertia veriflora, 161 glottis, 167 Padus, 149; melanocarpa, 149, 177 Philippine fungi, Bibliography and Palaeomyces, 75 new species of, 253 Palmoxylon cellulosum, 74, 77, 79 Pholiota, 316; aegerita, 113; candi-Panaeolus, 52, 69, 187; acidus, 187; cans, 53 campanulatus, 56, 298, 299; di-Phoma Lupini, 175 gressus, 187; epimyces, 52, 69; pa-Photographs and descriptions of cuppilionaceus, 317; reticulatus, 313; fungi-III. Peziza domiciliana and rufus, 313; variabilis, 313; vene-Peziza repanda, 195; IV. Peziza nenosus, 186 clypeata, 235 Panellus stypticus, 298, 299 Phragmidium imitans, 155; montiva-Panicum capillare, 127; fasciculatum, gum, 155; occidentale, 155; Peckianum, 155; Potentillae, 155; spe-Panus stipticus, 185 ciosum, 152

```
114; subelatinus, 218; subex-
Phragmites, 128; communis, 127
                                          cavatus, 219; subhaedinus, 218;
Phyllachora Dasylirii, 148; graminis,
                                          tremelliformis, 218
  148; Trifolii, 148
Phyllanthus grandifolius, 24
                                        Plicaria repanda, 198
Phylloporus rhodoxanthus, 298
                                        Plowrightia morbosa, 149; neomexi-
Phyllosticta Atriplicis, 176; baccha-
                                          cana, 149
                                        Pluteus cervinus, 313
  ridis, 55; cruenta, 176; maurandiae,
  55; medeolae, 55; oakesiae, 55;
                                        Poa pratensis, 147
  pachysandrae, 55; raui, 55; rhex-
                                        Poaceae, 16, 19, 21
  iae, 55; Symphoricarpi, 176
                                        Podosphaera Oxycanthae, 149
Physalacria inflata, 295
                                        Podostemaceae, 19
Physalis comata, 151; neomexicana,
                                        Podostroma alutacea, 295
  169
                                        Pogonomyces hydnoides, 56, 314
Physalospora aurantia, 149
                                        Poinciana pulcherrima, 18
Physarella oblonga, 36
                                        Poinsettia dentata, 168
Physarum bogoriense, 206; brunneo-
                                        Polemonium molle, 147; reptans, 137
  lum, 35; caespitosum, 204; chryso-
                                        Polygonum baxiforme, 147, 176; puc-
                                          tata, 43; ramosissimum, 147
  trichum, 205; citrinellum, 204; co-
  lumbinum, 201, 213; compactum,
                                        Polypodium, 44
  200, 201, 213; compressum, 202;
                                        Polyporaceae, 250, 296, 315
  cupriceps, 205; cupripes, 205; de-
                                        Polypores, A new genus of resupi-
  cipiens, 206; dictyospermum, 35;
                                          nate, 56
  didermoides, 36, 204; flavicomum,
                                        Polyporus, 15; admirabilis, 7-12, 15;
  205; globuliferum, 35, 203, 205;
                                          amorphus, 52; ampliporus, 215;
  lateritium, 36, 204; melleum, 200;
                                          arcularius, 173; argentatus, 215;
  murinum, 203; nutans, 36; pene-
                                          Berkeleyi, 250; cinnabarinus, 251;
  trale, 35; Petersii, 205; polycepha-
                                          cinnamomeus, 251; Curtisii, 251;
  lum, 35, 201; polycephalum var. obrusseum, 201, 202, 203; pulcher-
                                          destructor, 185; dichrous, 52; ele-
                                          gans, 296; Ellisianus, 178; fagi-
  rimum, 203, 209, 213; pulcherripes,
                                          cola, 296; flaccidus, 215; floccosus,
  203, 205, 206; pusillum, 203; rubi-
                                          215; gilvus, 251; hirsutus, 251;
  ginosum, 36; sinuosum, 206; stra-
                                          hispidus, 113; igniarius var. scaber,
  minipes, 36; vermiculare, 206; vi-
                                          214; lenziteus, 216; meyenii, 215;
  rescens, 202, 204, 213; viride, 35
                                          pargamenus, 251; rimosus, 214;
Physopella Artocarpi, 17
                                          robiniophilus, 251; sanguineus, 215;
Physotheca Halstedii, 145
                                          semipileatus, 251; spumeus mali-
Phytophthora cactorum, 109; infes-
                                          colus, 14, 15; tenuis, 217; tulipi-
  tans, 145
                                          ferus, 251; Underwoodii, 8, 9, 11,
Picea abies, 105; canadensis, 92; en-
                                          15; varius, 9; versicolor, 5, 251;
  gelmanni, 92; Parryana, 154; ru-
                                          xanthopus, 216; xanthosporus, 57
  bens, 92; sitchensis, 92
                                        Polystictus, 214; acutus, 215; affinis,
Pilosace, 69; algeriensis, 65, 69-71
                                          216; bogoriensis, 214; cinnabari-
Pileolaria patzcuarensis, 156
                                          nus, 215; crassipes, 216; cupro-
Pinus contorta, 91;
                      echinata, 91;
                                          nitens, 216; flabelliformis, 216;
  edulis, 309, 311; excelsa, 222; lam-
                                          floccosus, 215; luteus, 216; meye-
  bertiana, 91; monticola, 91; palu-
                                          nii, 215; nephalodes, 216; perula,
  stris, 91; ponderosa, 91, 178; pun-
                                          216; pterygodes, 216; saccatus,
  gens, 92; resinosa, 91; rigida, 91;
                                          216; sanguineus, 215; xanthopus,
  strobus, 92
                                          216
Pinus ponderosa, Note on western
                                        Polythelis Thalictri, 156
  red rot in, 178
                                        Polythrincium Trifolii, 177
Piper, 23; hispidum, 42
                                        Polytrichum commune, 294
Piperaceae, 23
                                        Populus aurea, 153; bolleana, 301;
Piptandenia macrocarpa, 20; pere-
                                          deltoides, 300, 306; nigra, 300,301,
  grina, 20
                                          305; virginiana, 301; Wislizeni, 154
Pisum sativum, 147, 148, 176
                                        Poronidulus conchifer, 296
Placosphaeria celtidis, 55; cornicola,
                                        Porotheleum, 56
  101
                                        Porotheliaceae, 56
Pleurotus albescens, 218; badius,
  219; dimidiatus, 298; Eryngii, 114;
                                        Porothelium, 56
  fulvifibrillosus, 219; geophilus,
                                        Porto
                                               Rican
                                                       parasitic
                                                                   fungi-II,
  218; petaloides, 252; phosphoreus,
                                          Studies in, 42
```

Porto Rico based on collections by F. L. Stevens, Uredinales of, 16 Potentilla monspeliensis, 155; strigosa, 155 Producing heart-rot of apple trees, Fungi, 5 Prospodium bahamense, 18; plagiopus, 18 Prunulus, 218; Abramsii, 220; adirondackensis, 220; alcaliniformis, 220; argillaceus, 220; atribrunneus, 220; atridiscus, 220; aurantiacus, 220; aurantiidiscus, 220; avellaneigriseus, 220; avellaneous, 220; brevipes, 220; caesiialbus, 220; carbonicola, 220; cervinialbus, 220; cinchonensis, 220; cinereiavellaneus, 220; collybiiformis, 220; cyaneobasis, 298; farinaceus, 220; flavicitrinus, 220; fuliginosus, 220; fumosiavellaneus, 220; fusipes, 220; gracillipes, 220; Grantii, 220; latericius, 220; Leaianus, 298; lepiotiformis, 220; leucophaeus, 220; longipes, 220; ludovicianus, 220; magnus, 220; margarita, 220; melleidiscus, 221; minutissimus, 221; murinus, 221; myceliosus, 221; niveipes, 221; occidentalis, ochraceicinereus, 221; paludicola, 221; parvulus, 221; pectinatus, 221; plumbeibrunneus, 221; pubescens, 221; roseipallens, 221; roseolus, 221; rutilantiformis, 221; Sabali, 221; scabripes, 221; subfumosus, 221; subpulverulentus, 221; subtenuipes, 221; syringeus, 221; tenuiculus, 221; testaceus, 221; trojanus, 221; viridigriseus, 221 Psathyrella disseminata, 54, 56, 252; impatiens, 54 Pseudocymopterus montanus, 161 Pseudomonas Citri, 112 Pseudopeziza Medicaginis, 150 Pseudotsuga taxifolia, 92 Psilopezia juruensis, 236, 237; nummularia, 235–238 Psilopeziza orbicularis, 237 Psoralea micrantha, 168 Ptelea, 136; trifoliata, 136, 141 Pteridium aquilinum, 165 Puccinia, 137; Absinthii, 156; Agropyri, 128, 129, 132, 133, 138, 139, 141, 157; alternans, 132, 141, 157; amphigena, 126, 127; angustata, 131, 132, 181; anomala, 139; apocrypta, 138; Artemisiae, 156: Aristidae, 156; Asparagi, 156; Asperifolii, 133, 141; atro-fusca, 156; canaliculata, 25, 157; Cannae, 25; Caricis-Asteris, 158; Caricis-Soli-

daginis, 158; Cenchri, 25; Cirsii, 157; Cladii, 19; Clematidis, 157; Cordiae, 17; Cressae, 157; Cryptandri, 157; Cynodontis, 20; Del-phinii, 157; Distichlidis, 136, 137, 141; Dulichii, 130, 140; emaculata, 127; Eleocharidis, 25; Ellisiana, 158; Epilobii-tetragoni, 158; Eriophori, 131, 132, 141; exitiosa, 18; extensicola, 130, 140, 158; farinacea, 24; Gayophyti, 158; graminis, 161, 181; Grindeliae, 158; Grosslulariae, 130, 141, 158; Helianthi, 158; hemispherica, 159; Huberi, 25; intermixta, 159; Jamesiana, 134, 159; Kelseyi, 136; Koeleriae, 159; lateritia, 25; levis, 25; lobata, 159; luxurians, 160; Magnusiana, 128; Menthae, 160; monoica, 160; montanensis, 137, 138, 139, 141, 160; Muhlenbergiae, 160; obliterata, 157; obtecta, 161; opulenta, 19; Oxalidis, 19; Philibertiae, 161; plagiopus, 18; poculiformis, 161; Prunispinosae, 16, 164; Pseudocymopteri, 161; punctata, 161; purpurea, 16; Rhamni, 128; rubigoverae, 129; Scirpi, 16; Seymouriana, 125, 130, 134, 135, 141; Sherardiana, 162; simillima, 127, 128; simplex, 139; Sorghi, 162; Spegazzinii, 19; Sphaeralceae, 162; Spilanthis, 19; splendens, 162; Stipae, 162; subnitens, 135, 141, 162; substriata, 25; Syndrellae, 25; Taraxaci, 163; tosta, 160; transformans, 18; tuberculans, 163; tumidipes, 136, 141, 163; uniporula, 130, 141; universalis, 163; Veratri, 163; Violae, 163; Windsoriae, 135, 141; Xanthii, 17, 164 Pucciniastrum Agrimoniae, 164; Myrtilli, 164; pustulatum, 164 Pulsatilla hirsutissima, 171 Pustularia Stevensoniana, 198 Pycnoporus cinnabarinus, 296; sanguineus, 56, 215 Pyrenomycetes, 98 Pyrola asarifolia, 154; chlorantha, 154; elliptica, 154; secunda, 154 Pyropolyporus conchatus, 296; Earlei, 174; igniarius, 174, 296; praerimosus, 174 Pythium debaryanum, 313

Quercus, 176, 177, 315; undulata, 174

Ramularia decipiens, 177; delphinii, 55; Grindeliae, 177; Lapsanae, 107; sambucina, 177

Ranunculus, 128; acris, 127, 128; bulbosus, 128: sceleratus, 127, 128

176;

busta, 299; uncialis, 299; vires-

cens, 122, 192

Ratibida columnifera pulcherrima, Rynchospora, 225 176 Ravenelia cassiaecola, 20; Cebil, 20; Saccharum officinalis, 42 Humphreyana, 18; mesilliana, 164; Salix, 153; amygdaloides, 153; Bebbiana, 153; exigua, 154; Fendlerpapillifera, 18; portoricensis, 16; versatilis, 164 iana, 154; irrorata, 154; lasiandra, Red rot in Pinus ponderosa, Note on 154; nigra, 154; petrophila, 154; Watsoni, 154; Wrightii, 153 western, 178 Remarkable fossil fungi, 73 Salsola pestifer, 162 Salvia coccinea, 24 Resupinate polypores, A new genus Sambucus mexicana, 177; microboof, 56 Resupinatus subbarbatus, 56 trys, 177 Saprolegnia Kauffmaniana, 108 Reticularia Lycoperdon, 40 Review of Kniep's Beiträge zur Sauvagesia erecta, 23 Schizophyllum commune, 5 Kenntnis der Hymenomyceten, III, Schizophyllus alneus, 56, 299 Schmaltzia Emoryi, 156; schmide-Rhamnus cathartica, 128; Purshiana, 128 lioides, 156 Schroeteriaster fenestrala, 24 Rhizoctonia, 314; Crocorum, 314; So-Scirpus, 131; americanus, 161; fluvilani, 314 Rhizopus nigricans, 313 atilis, 129; lacustris, 16 Rhus diversiloba, 106; Toxicoden-Scitamineae, 19 Scleria canescens, 24 dron, 106 Scleroderma aurantium, 299; vul-Ribes, 311, 314; Cynosbati, 130, 141; floridum, 130; grossularia, 311; inebrians, 176, 311; leptanthum, 309, 311; longiflorum, 130; longigare, 315 Sclerotites brandonianus, 75 Sclerotium Opuntiarum, 224 Scolochloa festucacea, 128 folium, 309, 311; mescalerium, 311; Scutiger Whiteae, 53 pumilum, 311; purpusii; 311; sanguineum, 104; valicola, 311; Wolfii, Seaver, Fred J., Fink's Ascomycetes of Ohio, 57; North American spe-151 Rigidoporus surinamensis, 56, 314 cies of Ascodesmis, 1; Photographs Rinodina angelica, 245 and descriptions of cup-fungi-III. Peziza domiciliana and Peziza re-Rosa, 43, 100, 155; Fendleri, 152, panda, 195, IV. Peziza clypeata, 235; The earth-inhabiting species 155; Maximiliani, 155; pecosensis, of Ascobolus, 93 Rosen, H. R., A known species of Sebacina, 316; atrata, 316; chlorassmut on a new host, 225 Rostkovites granulatus, 53 cens, 316; cinnamomea, 316; monticola, 316; plumbea, 316; Shearii, Rot in Pinus ponderosa, Note on western red, 178 316 Rubacer parviflorum, 155 Secale cereale, 141 Rudbeckia laciniata, 168 Senecio, 131; aureus, 132, 141; cacaliaefolius, 131; ductoris, 132; pa-Rubiaceae, 16 Rubus arizonicus, 155 luster, 131 Septobasidium laxum, 110 Rumex crispus, 177 Septogloeum, 106 Russula, 121, 122, 183, 299, 312; bi-Septoria, 106; adenocaulonis, 103; fida, 298; brunneola, 298; chamaeangularis, 103; Chrysanthemi, leontina, 298; compacta, 123; crus-104; darlingtoniae, 55; emaculata, tosa, 252; delica, 121, 124; emetica, 176; erythraceae, 55; Lapparum, 103; lupincola, 103; Lupini, 104; 298, 299; flava, 122, 183, 298; flavida, 122; foetens, 252, 299; Macrosporia, 104; Pisi, 176; foetentula, 315; fragiliformis, Polygonorum, 176; Ribis, 312; furcata, 123; granulata, 299; Rudbeckiae, 176; sanguinea, 104; lactea, 252; lutea, 122; Mariae, tinctoria, 55; Toxicodendri, 105 299; nigricans, 124; obscura, 123; Sequoia washingtoniana, 91 ochroleuca, 299; pectinata, 299; Sida, 44; hederacea, 161, 162 stricta, 299; subfragilis, 312; su-Simblum sphaerocephalum, 56

Sitanion, 138; elymoides, 138 Sium cicutaefolium, 129 Smilax, 127; hispida, 127 Smut on a new host, A known species of, 225 Solanaceae, 150 Solanum tuberosum, 145 Solidago, 130; canadensis, 103, 130, 140; fumosa, 103; juncea, 99; latifolia, 103; Pitcheri, 158, 162 Some Kentucky fungi, 249 Sophia incisa, 144, 145, 160, 162; ochroleuca, 145 Sophora sericea, 166 Sorghum halapense, 16; officinarum, 16; vulgare, 16 Sorosporium contortum, 170 South America, Myxomycetes from, Spartina, 125, 128, 134, 137; cynosuroides, 134, 141; gracilis, 136; Michauxiana, 134, 141 Species of Ascobolus, The earthinhabiting, 93 Species of Ascodesmis, North American, I Species of fungi, New or noteworthy, Species of smut on a new host, A known, 225 Spermacoce riparia, 25 Sphaeralcea angustifolia, 161; arenaria, 162; Fendleri, 162; incana, 162; lobata, 161, 162; marginata, 162 Sphaeriales, 145 Sphaeropsidales, 175 Sphaeropsis lineata, 101 Spilanthes oleracea, 19 Spiraea Menziesii, 98, 100, 102-104 Spondias mombin, 239, 242 Spongipellis fissilis, 14, 15; galactinus, 14, 15 Sporobolus airoides, 160; asperifolius, 160, 170; cryptandrus, 157, 172; utilis, 160; virginicus, 24 Spumaria licheniformis, 204 Stachys scopulorum, 147 Stagonospora Physocarpi, 103; Spiraeae, 103 Standley, Paul C., Fungi of New Mexico, 142 Steironema, 136, 137; ciliatum, 137, 141; Michauxiana, 137 Stemonitis dictyospora, 38, 39, 210, 213; digitata, 208; ferruginea, 39; fusca, 38, 208, 210; fusca var. dictyospora, 38; fusca var. rufescens, 38; fusca var. trechispora, 213; herbatica, 39; maxima, 208; porphyra, 208, 209, 213; splendens,

39; splendens var. Webberi, 39; tenerrima, 209; trechispora, 38, 39, 209, 213 Stemphylium nemopanthes, 107 Stenolobium Stans, 18 Stenorrhynchus lanceolatus, 21 Stereum bicolor, 56; elegans, 314; fasciatum, 250; lobatum, 56, 295; rugosum, 295; sericeum, 295 Stevens, Uredinales of Porto Rico based on collections by F. L., 16 Stipa Vaseyi, 172 Strobilomyces strobilaceus, 251, 296 Stropharia, 69; coprinophila, 69; semiglobata, 299; stercoraria, 174 Strophostyles, 165
Studies in Porto Rican parasitic fungi—II, 42
Sturgis, William C., Notes on the Myxomycetes of the Curtis Herbarium, 199; Myxomycetes from South America, 34 Sugar cane, Marasmius on, 115 Suillellus luridus, 297, 299 Sumstine, David R., A new species of Colus from Pennsylvania, 183 Symphoricarpos oreophilus, 176; rotundifolius, 148 Synchytrium fulgens, 144

Tapesia fusca, 295 Taraxacum Taraxacum, 163 Tecoma lepidota, 18; Leucoxylon, 18; Stans, 18 Teramnus uncinatus, 16, 45 Tests on the durability of American woods-I. Flask tests on conifers, Laboratory, 80 Tetranychus telarius, 107 Thalictrum, 133; dioicum, 132, 141; Fendleri, 132, 133, 156, 157 Thecaphora deformans, 170 Thelephora anthocephala, 250; magnispora, 53; palmata, 250; Schweinitzii, 250; vialis, 313 Thelephoraceae, 52, 250 Thuja occidentalis, 91; plicata, 91 Thyridium ceanothi, 55 Tidestromia lanuginosa, 144 Tilia, 315 Tilletia asperifolia, 170 Tilmadoche columbina, 200, 213 Tithymalus, 168; chamaesula, 151; luridus, 151 Tolysporella Nolinae, 170 Trametes acutus, 215; carnea, 251; cubensis, 314; limitata, 173; Pini, 178, 222 Tranzschelia punctata, 16, 164 Tremella albida, 250; lutescens, 295 Tremellaceae, 250

Tremellales, 295 Tremellodendron, 316; pallidum, 250; simplex, 316; tenue, 316 Tremellodon gelatinosum, 295 Trianthema Portulacastrum, 145 Trichamphora oblonga, 207 Trichia, 211; affinis, 41; Botrytis, 41, 211; Botrytis var. lateritia, 213; Botrytis var. munda, 41; decipiens, 41; favoginea, 41; persimilis, 41; serpula, 207; verrucosa, 41 Tricholoma imbricatum, 186 Trichosphaeria breviseta, 99 Trichothecium roseum, 177 Tricuspis seslerioides, 141 Tridens flavus, 135, 141 Trifolium Fendleri, 148; repens, 18, 148, 177; Rydbergii, 149 Triticum vulgare, 161 Tsuga canadensis, 91; heterophylla, 91; mertensiana, 91 Tubercularia vulgaris, 177 Tubifera ferruginosa, 40, 212; stipitata, 212, 213 Two parasitic mushrooms, 65 Tylopilus felleus, 297, 299; gracilis, Tylostoma, 56; fimbriatum, 175; gracile, 175; pedunculatum, 175 Tyromyces chioneus, 296; Ellisianus, 178; guttulatus, 296; semipileatus, 296

Ulmus americana, 102 Uncinula necator, 149; polychaeta, Uredinales, 150 Uredinales of Porto Rico based on collections by F. L. Stevens, 16 Uredineae in 1915, Cultures of, 125 Uredinopsis Pteridis, 165 Uredo Anthephorae, 19; Anthurii, 22; Artocarpi, 17, 26; Cordiae, 17; fenestrala, 24; globulosa, 22; Guacae, 21; Gymnogrammes, 25; gynandearum, 18; **Hameliae**, 23; Heliconiae, 19; Kaernbachii, 21; Olyrae, 21; Peperomiae, 23; Piperis, 23; pustulata, 21; Sauvagesiae, 23; superior, 17; venustula, 21: Wilsoni, 19 Urocystis Agropyri, 170; Anemones,

Uromyces, 136, 137, 139; albus, 165; appendiculatus, 165; Archerianus, 129, 165; Astragali, 165; Bidentis, 25; caryophilinus, 181; Cologaniae, 16; compactus, 165; Eriogoni, 166; Fabae, 166; Gentianae, 166; Glycyrrhizae, 166; Hordei, 139-141; Howei, 16; hyalinus, 166; ignobilis,

24; Limonii, 167; Medicaginis, 18, 167; mysticus, 167; occidentalis, 167; Pavoniae, 16; Peckianus, 129; plumbarius, 167; proeminens, 25, 167; Psoraleae, 168; Rudbeckiae, 168; Scirpi, 129; Scleriae, 241; Silphii, 168; Spartinae, 136; speciosus, 168; Trifolii, 18; Tranzschelii, 168

Uropyxis Amorphae, 168; Petalostemonis, 169; sanguinea, 169; Wootoniana, 169

Ustilaginales, 169

Ustilago Avenae, 171; bromivora, 171; Buchloes, 171; calcara, 171; Hieronymi, 171; Hilariae, 172; Hordei, 172; hypodytes, 172; levis, 172; Muhlenbergiae, 172; Zeae,

Vaccinium oreophilum, 164 Vaginata, 234; albocreata, 232; plumbea, 299; plumbea strangulata, 299 Vagnera amplexicaulis, 176 Venenarius, 231, 234; Frostianus, 299; glabriceps, 232, 233; muscarius, 299; pantherinus, phalloides, 123, 232, 233, 299; rubens, 299; solitarius, 231, 299 Venturia Dichiei, 149 Veratum speciosum, 163 Verbena Macdougalii, 147, 148 Vernonia albicaulis, 24 Verrucaria plumbaria, 245 Vicia americana, 165, 166 Viola canadensis, 163; nephrophylla, 163; pedatifida, 158 Volvaria bombycina, 65; hypopitys, 66; Loweiana, 65-68, 71; media, 67; parvula, 67; parvula var. major, 66; plumulosa, 66, 67, 71; pusilla, 67; umbonata, 67; villosa-volva, 67 Western red rot in Pinus ponderosa,

Note on, 178
Woods—I. Flask tests on conifers,
Laboratory tests on the durability
of American, 80

Xanthium commune, 164; longirostere, 17

Xanthoporia, 56; Andersoni, 57

Ximenesia exauriculata, 145, 148

Xylaria Hypoxylon, 295; polymorpha, 295

Xylometron sanguineum, 215

Young, Esther, Studies in Porto Rican parasitic fungi—II, 42 Yucca macrocarpa, 146

Zygophyllidium bilobatum, 168